


3WW/TN-80/003





THE CHEYENNE TORNADO: A METSAT CASE STUDY

JAMES C. ST JOHN, CAPTAIN, USAF



THIRD WEATHER WING
OFFUTT AFB, NEBRASKA 68113

MARCH 27, 1980

APPROVED FOR PUBLIC RELEASE:
DISTRIBUTION UNLIMITED

3d Weather Wing
Offutt AFB, Nebraska 68113

80 4 21 051

This report approved for public release. There is no objection to unlimited distribution of this report to the public at large, or by DDC to the National Technical Information Service (NTIS).

This technical report has been reviewed and is approved for publication.

FOR THE COMMANDER

Daniel R. GORNELL, Lt Col, USAF

Chief, Aerospace Sciences Division

Unclassified
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

ŀ	REPOR	RT DOCUMENTATION PAGE	READ INSTRUCTIONS BEFORE COMPLETING FORM							
t	REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER							
14	3 WW/TN-80/003	AD-A082 564								
	4. TITLE (and Subtitle)		5. TYPE OF REPORT & PERIOD COVERED							
1	The second secon									
16	The Cheyenne To	rnado; A METSAT Case Study								
4		······································	6. PERFORMING ORG. REPORT NUMBER							
ı	7. AUTHOR(s)		8. CONTRACT OR GRANT NUMBER(s,							
z. I	7. AUTHOR(S)									
101	JAMES C./ST. JO	HN∕ Capt								
\hookrightarrow		,								
ł	9. PERFORMING ORGANIZ	ATION NAME AND ADDRESS	10 PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS							
l	3d Weather Wing	, Aerospace Sciences	ARCH & WORK SHIP RESIDENCE							
		t AFB, Nebraska 68113								
l										
	11. CONTROLLING OFFIC	E NAME AND ADDRESS	12. REPORT DATE							
		-11 1 57	Mar ca 22, 20 80							
- 1	(Same as item 9)	LIA- NUMBER OF PAGES							
- 1	14 MONITORING ACENCY	NAME & ADDRESS(if different from Controlling Office)	15. SECURITY CLASS. (of this report)							
ı	14. MONITORING AGENCY	NAME & ADDRESS(II different from Controlling Office)	15. 5255							
- 1	(0 0	_	Unclassified							
- 1	(Same as item 9)	15a. DECLASSIFICATION DOWNGRADING							
ı			SCHEDULE							
ŀ	16. DISTRIBUTION STATEMENT (of this Report)									
	Approved for Public Release: distribution unlimited 17 DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)									
Ī										
ł	NA									
<u> </u>	18. SUPPLEMENTARY NOTES NA									
1										
J										
}										
1										
[19. KEY WORDS (Continue on reverse side if necessary and identify by block number)									
ĺ	METSAT TORNADO									
ı	Severe Convecti	ve Weather								
l	Meso-scale	vo moderior								
- 1	Visual Imagery		•							
t	20. ABSTRACT (Continue o	n reverse side if necessary and identity by block number)								
- 1	This Technical	Note describes the Meteorological S	Satellite (METSAT imagery, of							
)	the thunderstorm outbreak, which produced the Cheyenne, Wyoming tornado. It									
	discusses METSAT Imagery signatures for severe weather and the application of									
- 1	METSAT Imagery in providing weather support to certain Air Force operations.									
- 1										
ı										

DD 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE


Unclassified
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

ACKNOWLEDGEMENTS

I would like to thank CMSgt Eugene M. Weber for sharing his knowledge of the mechanics of producing technical publications. I would also like to thank Sgt Farley of 3 WW/DNS and TSgt Jarlett of the Offutt AFB photo lab for their help with the graphics in this study, and Major Henry A. King III for his editorial help. Special thanks go to Mrs. Diane Brooks for her typing skills.

TABLE OF CONTENTS

	Page
List of Illustrations	. ii
Introduction	. 1
Synoptic Situation	. 2
GOES Imagery	. 2
METSAT Imagery Tornado Signature	. 7
Need For GOES Imagery	. 7
Conclusion	. 7



LIST OF ILLUSTRATIONS

Figu	ıres												Page
1.	The Cheyen	ne Tornado .				•	•		•				1
2.	Damage to	ANG C-130					•					•	1
3.	SMS - 2 Vi	sual Imagery,	1646Z	16 July	1979	•							3
4.	SMS - 2 Vi	sual Imagery,	1746Z	16 July	1979					•		•	3
5.	SMS - 2 Vi	sual Imagery,	1846Z	16 July	1979	•							3
6.	SMS - 2 Vi	sual Imagery,	1946Z	16 July	1979			•				•	4
7.	SMS - 2 Vi	sual Imagery,	2016Z	16 July	1979				•			•	4
8.	SMS - 2 Vi	sual Imagery,	2046Z	16 July	1979	•				•			4
9.	SMS - 2 Vi	sual Imagery,	2116Z	16 July	1979								5
LO.	SMS - 2 Vi	sual Imagery,	2146Z	16 July	1979								5
ll.	SMS - 2 Vi	sual Imagery,	2216Z	16 July	1979				•		•	•	5
ι2.	SMS - 2 Vi	sual Imagery,	2246Z	16 July	1979		•	•	•				6
13.	SMS - 2 Vi	sual Imagery,	2346Z	16 July	1979								6




Figure 1. The Cheyenne Tornado (Photo courtes, Capt J. James, OL-A, 9 WS F.E. Warren AFB, WY).

1. Introduction.

At 2135Z on 16 July 1979 a tornado struck Cheyenne, Wyoming (Figure 1). The funnel cloud developed over Francis E. Warren AFB, proceeded along the northern boundary of the base and continued eastward into Cheyenne. It caused one fatality and over \$40 million damage in the civilian community (Ostby and Wilson, 1980). Damage to Air National Guard C-130s at the Cheyenne Airport totaled several million dollars, (Figure 2).

One problem in forecasting such an occurrence is the lack of surface observations and radar data in the sparsely populated regions of the west. Satellite data covers a continuum of space, and the resolution of GOES imagery allows mesoscale analysis. GOES satellite imagery is not dependent upon surface reporting stations or location of radars. These data are available at 30 minute intervals to units with GOES or WSFO taps. An analysis of the available GOES imagery from the afternoon of 16 July suggests a technique for predicting the occurrence of severe convective weather.



Figure 2. Wyoming ANG C-130 damaged by the Tornado (Photo courtesy Capt J. James OL-A, 9 WS, F. E. Warren AFB, WY).

2. Synoptic Situation.

There was a high potential for severe convective weather over the high plains of southeastern Wyoming and northeastern Colorado on 16 July 1979. A shortwave trough with weak to moderate PVA was evident at 500mb. A 90kt westerly jet at 200mb was over the area, providing a venting mechanism for deep convection. There was strong thickness advection in the low levels, as well as high moisture content. The atmosphere was conditionally unstable. Since no sounding was available over Cheyenne, the exact vertical thermal structure is unknown. Analysis of surrounding soundings, however, indicates that a layer of warm dry air overlaid a cool moist layer near the surface. This is a convectively unstable situation with the potential for explosive overturning. All the ingredients for severe weather were present.

Since we began forecasting severe weather we have faced this limitation: We can forecast areas of severe weather potential with considerable skill, but we are unable to forecast the time and place of occurrence, until the thunderstorms develop and can then be tracked by radar. We are unable to forecast tornado occurrence until one is sighted, or a radar signature is observed. Then extrapolation of movement is used to predict the path of the storm. Most base weather stations within Air Weather Service will not issue a warning for severe convective weather elements until their signatures are displayed upon a radar or until reported by observation. The warning lead time is dependent, therefore, on how far upstream in time and space the severe weather signatures or elements are observed. Therefore, on days such as July 16, even though there may be great potential for severe convective weather, point warnings are often not issued until the first occurrence of a severe weather element.

3. GOES Imagery.

At 1646Z the satellite imagery (Figure 3) showed the first cumulus development, near Casper, to the north of the Cheyenne area. The 1746Z GOES picture (Figure 4) showed the first thunderstorm developing southwest of Laramie. Some small cumulus can be seen developing over the Laramie Mountains between Laramie and Cheyenne. By 1846Z (Figure 5) a mature thunderstorm cell was located over Cheyenne with a second mature cell just to the west. Two small cells in the building stage were located to the north. The National Weather Service observation at the Cheyenne Airport included thunder from 1826Z to 1829Z. By 1946Z (Figure 6) a line of thunderstorm cells extended from just south of the Colorado border to the area just south of Laramie, then northeastward to Kimball, Nebraska. Cells were also developing north-to-scuth along the ridgeline of the easternmost ranges of the Rockies. A large cluster of these cells developed from the west-to-southwest of Laramie. This was the first indication of organization of the cells into mesoscale convective systems. Note also the two cloud clusters at point C and the small cell at point D.


The picture for 2016Z (Figure 7) shows the two large cells at C have remained nearly stationary but have become better organized. The small cell at point D has not grown appreciably but has moved rapidly to the northeast. The line to the scutheast has remained stationary. The area of cells to the southwest of Laramie continued to grow. A recapitulation of events at this point would indicate that mesoscale cloud systems have grown and become organized while remaining quasi-stationary, except for the somewhat intriguing movement of the one cell labelled D.

Figure 8 shows the situation at 2046Z. Cell D has developed dramatically into a mature thunderstorm. Cells Cl and C2 have remained stationary while the line of well. (A) have moved to the southeast. The cloud mass southwest of Laramie, new shows two large distinct cells Bl and B2. B2 is beginning to merge with Cl and C2. The variation in movement between the cells would seem to indicate the convective interaction has developed into a mesoscale circulation system which dominated the matir ment of northeastern Colorado, southeastern Wyoming and the Nebraska panhandle.

The 2116Z imagery (Figure 9) shows a dramatic departure. There are now four invitant cells in the region. Two have persisted for several hours and have obviously reached quasi-steady-state supercell status. These are cell D which has continued to move eastward and cell A which has grown "fatter" and has lost its line appearance. Tells B2, C1 and C2 have merged and are now labelled merely C. Cells A and B are beginning to merge, just to the west of Cheyenne. Practically the entire area between Casper, Wyoming and Denver, Colorado east to the Nebraska border is covered by these four thunderstorm clusters.

By 2146Z (Figure 10) cells A and B have fully merged and C and D have begun to merge. The merged A and B cells extend east and west of Cheyenne. Figures 11, 12 and 13 show that by 2246Z the cells all have merged into one large group.

Figure 3. SMS - 2 Visual imagery at 1646Z shows first cumulus development near Casper (A).







Figure 5. SMS - 2 Visual Imagery at 1846Z. A thunderstorm cell is over Cheyenne (A). There is considerable thunderstorm development over the Laramie mountains (B). Two small cells are beginning to develop north of Laramie (C).






Figure 6. SMS - 2 Visual imagery at 19462. The thunderstorms have organized into a line from the Colorado border to Kimball, Nebraska (A). Another cluster of cells is growing over the eastern slope of the Rockies (B). The two small cells (C) from the last picture have grown into cloud clusters. Another small cell has begun development (D).






Figure 7. CMS - 2 Visual imagery at 20160. The line of thunderstorms (A) has remained nearly stationary, as has the cluster of cells over the mountains (B). The two clusters (Cl and C2) have changed little in 30 minutes. Note, however, the rapid movement of cell D.







Figure 9. SMS - 2 Visual imagery at 21167. Cells B2, S1 and C2 have merged and are now designated B. The line A now has a circular appearance, and with cell D have reached super cell status. Cell A and cell B appear to be merging, just to the west of the Cheyenne



First 11. TWO - 3 Wishal imagery at 1.6%. Wello Cand Dehave negrots.



Who are 10. Two - Thomas Inches on 1800. The minutes after the tornais one amence, sells A and F have completely merged over the Theorems area. Tells O and I have been to merge.






Figure 12. CMS - 2 Visual imagery at 2246Z. The two large cells appear to be merging.



Figure 13. SMS - 2 Visual imagery at 23467. The entire area is covered with thunderstorm cirrus chields.

4. METSAT Imagery Tornado Signature.

The relationship between mercing cells on radar displays and the occurrence of tornadoes have been discussed (e.g. The Ft. Bucker Tornado (Miller, 1972)). A limitation of satellite imarry is that the precipitation cells cannot be located, only the cloud tops are visible. Even so, close analysis of these data indicate that two large thunderstorms merced over the Cheyenne area at the time of the tornado occurrence. Furthermore, the imagery indicates the interaction of the cells during the development of this large convective system. Furdom (1979) demonstrated the primary importance of convective scale interaction in the evolution of deep convection.

This indicates that careful analysis and met watch of these NOES data by a trained forecaster could have provided strong indication of severe thunderstorm activity more than an hour before the tornado occurrence. Furthermore, 19 minutes before tornado occurrence, there was indication of merging cells; a tornado "signature".

5. Need For IOES Imagery.

Radar data would provide a much more complete analysis of such occurrence. METSAT imagery is not a substitute for radar. Rather, the two systems are complementary. Purdom (1979) states:

"Rapid interval patellite imagery combined with radar data must be the basis for short term forecasting (noweasting) of a sweetien. This is because they are the only meteorological observing systems that can menitor the convective scale interactions that are so vital in the maintenance of deep convective activity".

For units which have no radar, yet are tasked to provide weather wersing suspert, a 10E3 Tap will so a 1 no way to "close the gap" in mesoscale analysis. Hiven this met watch tool, forecasters can closely monitor the data for signs of leveleging singular to restrict, all and system movement, and sinus of severity (rapid on a movement, high clouds tops from infrared imagency, morning cells, etc.). The stilling to forecast the time of communese and degree of severity of the severe a syestive weather elements will not be as spect as with a radar, as wellness will have to be rape seneral with a resultant increase of timing error that present intersity over r. This is still preferable to a degree of symming less time.

ϵ . Constant a..

A review of ICEA imageny from it does to be determined to itself a more instituted storm medianism was responsible for the Rependent means. This leads to the conciusion that mention will in METUAT imaging uncolled a solution of a tornado signature for severe weather forecasting runs seed. This can be writed by these evidence of the need for ICEA Tape at 10000 AMO for resting a list.

Millon, B. J., 1972: Notes on Analusis and Severe Storm Forecasting Procedures of the Air Force Global Weather Central, AMNUTE 1991.

Outby, W. P. and Wilson, L. V., 1980: Temade! Westerwise, e. a. gl-85.

Fundom, J. F. W. 1979: The Development and Evalution of Deep Convection, I results to I lumbs: Eleventh Confloration on Covers Local Starmer, pp. 186-18.